Work Activities Deemed Not Critical (PT)

Work Activity Statement	Rationale	
PATIENT/CLIENT ASSESSMENT		
Tests & Measures		
Muscle Performance		
Select and perform tests and measures ofelectrophysiological function using surface electrodes (e.g., surface EMG)	Survey responses suggest that this activity is trending down compared to the 2011 survey. The Task Force suggested this activity may be better classified under Therapeutic Modalities: Perform and/or train patient/client/caregiver in biofeedback therapy.	
electrophysiological function using needle insertion (e.g., nerve conduction)	The Task Force suggested this may not be an entry-leve activity.	
muscle integrity (e.g., ultrasound imaging)	While the survey responses suggest that this activity is trending up from the 2011 survey and is being covered more in entry-level curriculum, the Task Force felt that the evidence was not quite compelling enough to include this activity on the NPTE yet. The Task Force recommended to survey this activity again at the next practice analysis.	
Neuromotor Development & Sensory		
Integration		
Select and perform tests and measures oforal motor function, phonation, and speech production	The Task Force suggested that the oral motor function portion of this activity may be better classified under Systems Review: Perform screen of the gastrointestinal system. The Task Force felt there may be limited PT related tests and measures in this category and they may not be entry level.	
INTERVENTIONS		
Manual Therapy Techniques Perform dry needling	This activity was excluded based on the empirical decision rule of the survey responses and the agreement of the Task Force. The Task Force recommended to survey this activity again at the next practice analysis.	
Integumentary Repair & Protection Techniques		
Perform and/or train		
patient/client/caregiver in		
hyperbaric therapy	This activity was excluded based on the empirical decision rule of the survey responses and the agreement of the Task Force.	
negative pressure wound therapy	The Task Force questioned whether this was an entry- level activity and suggested the frequency of its use may depend on specific practice setting (e.g., hospital, military).	

Thoranoi	ıtic	Modalities	
inerabet	มเเต	iviodalities	

Perform and/or train patient/client/caregiver in... ...hydrotherapy using contrast baths/pools This activity was excluded based on the empirical decision rule of the survey responses and the agreement of the Task Force. ...phototherapy (laser light) This activity was excluded based on the empirical decision rule of the survey responses and the agreement of the Task Force. The Task Force suggested this activity is highly equipmentdependent (if the equipment is not available, the work activity will not be performed). ...monochromatic infrared agent procedures This activity was excluded based on the empirical (e.g., light emitting diodes [LEDs]) decision rule of the survey responses and the agreement of the Task Force. The Task Force suggested this activity is highly equipmentdependent (if the equipment is not available, the work activity will not be performed). ...diathermy This activity was excluded based on the empirical decision rule of the survey responses and the agreement of the Task Force. ...dry heat thermotherapy (e.g., Fluidotherapy) This activity was excluded based on the empirical decision rule of the survey responses and the agreement of the Task Force ...paraffin bath thermotherapy This activity was excluded based on the empirical decision rule of the survey responses and the agreement of the Task Force. The Task Force felt that practice is moving more toward screening for appropriateness of this thermal modality, but this activity could be performed in many settings other than rehabilitation. This activity was excluded based on the empirical ...shockwave therapy decision rule of the survey responses and the agreement of the Task Force.